Family list 7 family members for: WO2004011400 Derived from 6 applications.

- 1 METHOD OF DEUTERATING AROMATIC RING Publication info: AU2003248267 A1 - 2004-02-16
- 2 A METHOD OF DEUTERATION OF AN AROMATIC RING Publication info: CA2493773 A1 2004-02-05
- 3 Method of deuterating aromatic ring Publication info: CN1675145 A - 2005-09-28
- 4 METHOD OF DEUTERATING AROMATIC RING Publication info: EP1535889 A1 - 2005-06-01 EP1535889 A4 - 2006-04-05
- 5 METHOD OF DEUTERATING AROMATIC RING Publication info: RU2005105297 A - 2005-07-20
- 6 METHOD OF DEUTERATING AROMATIC RING Publication info: WO2004011400 A1 - 2004-02-05

Data supplied from the esp@cenet database - Worldwide

METHOD OF DEUTERAYING AROMATIC RING

Publication number: WO2004011400

Publication date:

2004-02-05

Inventor:

ITO NOBUHIRO (JP); MAESAWA TSUNEAKI (JP);

MUTO KAZUSHIGE (JP); HIROTA KOSAKU (JP);

SAJIKI HIRONAO (JP)

Applicant:

WAKO PURE CHEM IND LTD (JP); ITO NOBUHIRO (JP); MAESAWA TSUNEAKI (JP); MUTO KAZUSHIGE (JP); HIROTA KOSAKU (JP); SAJIKI HIRONAO (JP)

Classification:

- international:

C07B59/00; C07C39/04; C07C39/28; C07C51/00; C07C51/347; C07C209/68; C07C315/04; C07C211/46; C07C211/51; C07C317/14; C07B59/00; C07C39/00; C07C51/00; C07C51/347; C07C209/00; C07C315/00; C07C211/00; C07C317/00; (IPC1-7): C07B59/00; C07C5/00; C07C15/16; C07C37/00; C07C39/04;

C07C39/28; C07C51/347; C07C63/06; C07C63/08; C07C209/68; C07C211/46; C07C217/84; C07C315/04;

C07C317/14; C07M5/00

- european:

C07B59/00; C07B59/00C; C07C5/00; C07C39/04;

C07C39/28; C07C51/00; C07C51/347; C07C209/68;

C07C315/04

Application number: WO2003JP08783 20030710 Priority number(s): JP20020219005 20020726

Also published as:

EP1535889 (A1) CN1675145 (A) CA2493773 (A1) AU2003248267 (A

Cited documents:



JP63198638 JP63030450 JP63030435 JP47012567 XP004017175 more >>

Report a data error he

Abstract of WO2004011400

A method of deuterating a compound having an aromatic ring in the presence of an activated catalyst. In particular, a method of deuterating a compound having an aromatic ring characterized in that a compour having an aromatic ring is reacted with a deuterium source in the presence of an activated catalyst selected from among a platinum catalyst, a rhodium catalyst, a ruthenium catalyst, a nickel catalyst and a cobalt catalyst.

Data supplied from the esp@cenet database - Worldwide